SNAPSHOOT
ENGINEERED FOR A SAFER TOMORROW

현장을 이해하고, 필요한 솔루션을 연결합니다.

산업안전 · 추락방지 시스템
산업용 소재 · 건축자재
제품 제안부터 공급·시공·설치·교육까지



COMPANY INFORMATION

스냅슛 (SNAPSHOOT)

ADDRESS (15431) 경기도 안산시 단원구 풍전로 37-9 트리플렉스 301동 118호
TEL 0507-1374-9229
FAX 0504-481-6749
E-MAIL bkkim@snapshoot.co.kr

TECHNICAL DOCUMENTS

인증서 · 데이터시트 · 기술자료

제품의 규격과 성능, 인증 및 적용 조건을 확인하기 위한 자료입니다.

ABOUT SNAPSHOOT

제품 공급을 넘어, 현장에 맞는 해결책으로

스냅슛은 산업현장의 작업 환경과 요구 조건을 바탕으로
적합한 제품과 시스템을 제안합니다. 현장 실사를 통해 설치 조건을 확인하고,
공급부터 시공·설치 및 사용 교육까지 필요한 과정을 함께합니다.

추락방지 시스템

안전대부착설비 · 수평·수직 라이프라인
레일 시스템 · 리지드 포스트

밀폐공간 진입·구조 솔루션

데빗암 · 삼각대 · 구조용 윈치
윈치 안전블록 · 구조장비

산업안전 보호구

전신형 안전대 · 랜야드 · 안전블록
작업 환경에 맞춘 개인보호구

산업용 소재·건축자재

산업용 테이프 · 접착제 · 기능성 소재
구조용 스크류 · 목조건축 자재

SNAPSHOOT
ENGINEERED FOR A SAFER TOMORROW

홈페이지 바로가기 →
문서 라이브러리 바로가기 →
시공사례 바로가기 →
문의하기 →

본 페이지는 스냅슛의 자료 제공 안내용 표지입니다. 첨부된 인증서 및 기술자료의 발행 주체, 인증 대상과 적용 범위는 해당 원문을 기준으로 확인해 주세요.

NEOGEN® Culture Media

EN (English)

Product Instructions

Half Fraser (Demi-Fraser) Broth (ISO)

Intended Use

Half Fraser (Demi-Fraser) Broth (ISO) is a selective medium used for the primary enrichment of *Listeria monocytogenes* and *Listeria* spp., as described in ISO 11290–1:2017. Half Fraser (Demi-Fraser) Broth (ISO) is not intended for use in the diagnosis of disease or other conditions in humans.

Description

Half Fraser (Demi-Fraser) Broth (ISO) was developed as a modification of UVM medium and is based on the formula as described by Fraser and Sperber. *Listeria* spp. hydrolyze the esculin to esculetin which reacts with the ferric ammonium citrate resulting in a black precipitate and a visible positive reaction. Lithium chloride suppresses the growth of *Enterococci* in the medium, as *Enterococci* can also hydrolyze esculin. Enzymatic digest of animal tissues, casein, meat extract and yeast extract provide the essential vitamins, minerals, amino acids, nitrogen and carbon, while sodium chloride maintains the osmotic balance. The selective components acriflavine and nalidixic acid are blended into the base powder and inhibit the growth of *Escherichia coli*. Ferric ammonium citrate is added to the tempered broth after sterilization. According to ISO 11290–1:2017, the test portion is inoculated into the primary enrichment medium Half Fraser (Demi-Fraser) Broth (ISO), followed by secondary enrichment in Fraser Broth (ISO) and then sub-cultured onto Harlequin *Listeria* Chromogenic Agar according to Ottaviani & Agosti (700004790, 700004791, 700004792, 700004793, 700004794, NCM1004). The medium conforms to the performance and formulation requirements of ISO 11290–1:2017.

Typical Formulation	
Enzymatic Digest of Animal Tissues	5.0 g/L
Enzymatic Digest of Casein	5.0 g/L
Meat Extract	5.0 g/L
Yeast Extract	5.0 g/L
Esculin	1.0 g/L
Disodium Hydrogen Phosphate	9.6 g/L*
Potassium Dihydrogen Phosphate	1.35 g/L
Sodium Chloride	20.0 g/L
Lithium Chloride	3.0 g/L
Acriflavine	0.0125 g/L
Nalidixic Acid	0.01 g/L

pH: 7.2 ± 0.2 at 25°C

*Equivalent to Disodium Hydrogen Phosphate Dihydrate 12.0 g/L

Formula is adjusted and/or supplemented as required to meet performance specifications.

Supplement	
700004872 | NCM4009-0.5 or 700002053 | BP0220010	Ferric Ammonium Citrate or Fraser Broth Supplement

Precaution

Refer to safety data sheets.

Dehydrated Culture Medium Preparation

1. Dissolve 55 g of the medium in one liter of purified water.
2. Mix thoroughly.
3. Autoclave at 121°C for 15 minutes
4. Cool to 45–50°C.
5. Aseptically add 2 vials of 700004872 | NCM4009–0.5* Ferric Ammonium Citrate Supplement or 1 vial of 700002053| BP0220010 Fraser Broth Supplement and mix thoroughly before dispensing.

*Larger vials may be available. Please see appropriate supplement data sheet for availability and preparation instructions.

Ready-to-Reconstitute Media Bag Preparation

Please refer to the Instructions for Use for the range.

Test Procedure

To isolate *Listeria monocytogenes* and other *Listeria* spp., refer to ISO 11290–1:2017.

Quality Control Specifications

Dehydrated Appearance: Powder is homogeneous, free flowing, and beige.

Prepared Appearance: Prepared medium is yellow to amber and clear with none to slight precipitate.

Expected Cultural Response: The medium was prepared according to label directions, incorporating 700004872 | NCM4009–0.5* Ferric Ammonium Citrate Supplement or 700002053 | BP0220010 Fraser Broth Supplement and 10 mL volumes inoculated with the organisms listed below. Cultures were incubated at 30 ± 1°C under aerobic atmosphere and examined for growth at 22–26 hours. Following incubation, 10 µL was sub-cultured onto TSA at 37 ± 1°C and examined for growth at 22–26 hours or onto Harlequin® *Listeria* Chromogenic Agar (700004790, 700004791, 700004792, 700004793, 700004794, NCM1004)* at 37 ± 1°C and plates examined for growth at 40–48 hours.

Microorganism	Wdcm	Approx. Inoculum (CFU)	Expected Results				Actual Results
			Growth in HF	Reaction in HF*	Recovery on HLCA**	Recovery on TSA	
Listeria monocytogenes + *Escherichia coli* + *Enterococcus faecalis*	00021 00013 00087	10 – 100 > 10^4 > 10^4	Growth	Black coloration	>10 cfu Blue colonies with opaque halo	Not applicable	Meets Expected Result
Listeria monocytogenes + *Escherichia coli* + *Enterococcus faecalis*	00109 00012 00009	10 – 100 > 10^4 > 10^4	Growth	Black coloration	>10 cfu Blue colonies with opaque halo	Not applicable	Meets Expected Result
Escherichia coli	00012	> 10^4	Inhibition	None	Not applicable	Total inhibition	Meets Expected Result
Escherichia coli	00013	> 10^4	Inhibition	None	Not applicable	Total inhibition	Meets Expected Result
Enterococcus faecalis	00009	> 10^4	Partial to Complete Inhibition	None	Not applicable	< 100 cfu	Meets Expected Result
Enterococcus faecalis	00087	> 10^4	Partial to Complete Inhibition	None	Not applicable	< 100 cfu	Meets Expected Result

The organisms listed are the minimum that should be used for quality control testing.

Results

Listeria is presumptively indicated by the blackening of Half Fraser (Demi-Fraser) Broth (ISO) after 25 ± 1 hours incubation at 30°C ± 1°C.

*Sub-culture onto LCA provides presumptive *L. monocytogenes* or *L. ivanovii*, if colonies are blue-green surrounded by an opaque halo. Consider presumptive *Listeria* spp. if blue-green colonies with or without opaque halo.

Expiration

Refer to expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if appearance has changed from the original color. Expiry applies to medium in its intact container.

Limitations of the Procedure

1. Some strains of *L. monocytogenes* exposed to stress conditions, particularly acid stress, can show a very weak halo.
2. Some rare *L. monocytogenes* are characterized by a slow PIPLC activity. Such organisms will take longer to develop a halo.

Storage

Store dehydrated culture media at 2–30°C away from direct sunlight. Once opened and recapped, place the container in a low humidity environment at the same storage temperature. Protect from moisture and light by keeping container tightly closed.

Store ready-to-reconstitute media bags (NCM3205) at 10–25°C away from direct sunlight. Once reconstituted store at 10–25°C away from direct sunlight for up to 5 days (providing asepsis is maintained).

References

1. Fraser, J.A. and Sperber, W.H. (1988). Rapid detection of *Listeria* spp in food and environmental samples by esculin hydrolysis. J. Food Protect. 51, No.10, 762-765.
2. McClain, D. and Lee, W.H. (1989). FSIS method for isolation of L. monocytogenes from processed meat and poultry products. Lab.Comm.No.57, Revised May 24, (1989). US Dept of Agric. FSIS, Microbiol. Div.
3. ISO 11290-1:2017 Microbiology of the food chain- Horizontal method for the detection and enumeration of *Listeria monocytogenes* and *Listeria* spp.- Part 1: Detection method
4. ISO 11290-2:2017 Microbiology of the food chain- Horizontal method for the detection and enumeration of *Listeria monocytogenes* and *Listeria* spp. - Part 2: Enumeration method
5. ISO 11133:2014+A1:2018 Microbiology of food, animal feed and water - Preparation, production, storage and performance testing of culture media.

NEOGEN

Neogen Corporation, 620 Lesher Place, Lansing, MI 48912 USA.
© Neogen Corporation 2026. All rights reserved. Neogen is a registered trademark of Neogen Corporation.
All other marks mentioned are the property of their respective owners.